

Supplementary Table 2. Clinical information of suicide attempters and Bonferroni post-hoc comparisons

Variable	Adult group (n = 656)	Elderly group (n = 123)	χ^2 or Fisher exact test	p-value
Past psychiatric treatment			11.946	0.018
Likely to have had psychiatric problems (never seen a doctor)	201 (37.5)	57 (50.0)		
Likely to have not had psychiatric problems (never seen a doctor)	47 (8.8)	14 (12.3)		
Discontinued treatment	58 (10.8)	13 (11.4)		
Currently in treatment (with medication)	227 (42.4)	29 (25.4)		
Currently in treatment (no medication)	3 (0.6)	1 (0.9)		
Past suicide attempt history			7.398	0.007
Yes	190 (33.9)	23 (20.7)		
No	371 (66.1)	88 (79.3)		
Physical treatment after past suicide attempts			1.097	0.295
Visited hospital	107 (68.6)	16 (80.0)		
Did not visit hospital	49 (31.4)	4 (20.0)		
Psychiatric treatment after past suicide attempts			1.975	0.372
Received treatment	91 (60.3)	11 (57.9)		
Not received treatment (recommended)	44 (29.1)	4 (21.1)		
Not recommend treatment	16 (10.6)	4 (21.1)		
Family history of psychiatry			0.072	0.789
Yes	355 (83.1)	70 (84.3)		
No	72 (16.9)	13 (15.7)		
Family history of suicide attempt			0.405	0.524
Yes	43 (10.5)	6 (8.1)		
No	365 (89.5)	68 (91.9)		
Awareness of suicide			10.334	0.006
Admit	84 (16.0)	10 (10.5)		
Deny	430 (82.1)	78 (82.1)		
Uncheckable	10 (1.9)	7 (7.4)		
Past medical history			177.934	<0.001
No underlying disease	404 (72.0)	15 (12.6)		
Recent acute disease	9 (1.6)	7 (5.9)		
Chronic disease (not interfering with daily life)	125 (22.3)	58 (48.7)		
Chronic disease (interfering with daily life)	23 (4.1)	39 (32.8)		
Past psychiatric admission count ^{a)}	2.01	0.15	13.120	<0.001
Past suicide attempt count ^{a)}	0.85	0.35	356.332	<0.001

Values are presented as numbers (%) or mean value only.

^{a)}t-test.