

**Supplementary Table 1.** Distribution of departments in urine VRE diagnosis

Characteristic	Doxycycline treatment (n = 21)	Non-treatment (n = 62)	Total (n = 83)	p-value
Department of VRE diagnosis				0.324
Neurosurgery	3 (14.3)	10 (12.9)	13 (15.7)	
Nephrology	3 (14.3)	8 (12.9)	11 (13.3)	
Infection	6 (28.6)	4 (6.5)	10 (12.0)	
General surgery	1 (4.8)	8 (12.9)	9 (10.8)	
Hemato-oncology	1 (4.8)	7 (11.3)	8 (9.6)	
Gastroenterology	1 (4.8)	6 (9.7)	7 (8.4)	
Rehabilitation	1 (4.8)	6 (9.7)	7 (8.4)	
Rheumatology	1 (4.8)	3 (4.8)	4 (4.8)	
Endocrinology	1 (4.8)	2 (3.2)	3 (3.6)	
Urology	1 (4.8)	1 (1.6)	2 (2.4)	
Neurology	0	2 (3.2)	2 (2.4)	
Vascular surgery	0	1 (3.2)	2 (2.4)	
Plastic surgery	1 (4.8)	0	1 (1.2)	
Breast & thyroid surgery	1 (4.8)	0	1 (1.2)	
Cardiology	0	1 (1.6)	1 (1.2)	
Emergency	0	1 (1.6)	1 (1.2)	
Pulmonary	0	1 (1.6)	1 (1.2)	

Values are presented as number (%). VRE, vancomycin-resistant *Enterococcus*.